




MEDICO RESEARCH CHRONICLES

ISSN NO. 2394-3971

DOI No. 10.26838/MEDRECH.2021.8.3.498

Contents available at www.medrech.com

CLINICAL AND USG FINDINGS OF PATIENTS PRESENTED WITH BREAST PAIN

H. N. Ashikur Rahaman¹, Shravana Kumar Chinnikatti²

1. Registrar, Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, Dhaka, Bangladesh

2. Senior Consultant, Dept. Of Clinical Oncology, Enam Medical College Hospital, Savar, Dhaka, Bangladesh

ARTICLE INFO

Article History

Received: April 2021

Accepted: June 2021

Keywords: Breast Pain, Breast Cancer, Palpable Mass, USG Findings, Axillary Lymph Nodes.

ABSTRACT

Introduction: Breast pain also known as mastalgia is the dull ache in the breast, the cause is multifactorial. It can affect any age group. Breast cancer is the most common site-specific cancer in women and is the leading cause of death from cancer for women aged 20-59 years worldwide. In India, breast cancer has ranked number one cancer among females, with an age-adjusted incidence as high as 25.8 per 100,000 women and mortality of 12.7 per 100,000 women. **Objective:** To find out the Clinical and USG Findings of Patients Presented with Breast Pain. **Settings and Design:** Prospective cross-sectional descriptive study. **Materials and Methods:** Patients with complaints of breast pain in the age group 15 to 60 years with clinically no palpable mass attending to the Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, Dhaka, Bangladesh between January 2020 to December 2020 were included in the study. Detailed history with particular reference to age, duration of symptoms, Size, menstruation, marital status, parity, lactation, nipple discharge, and tenderness are recorded. Pain intensity was evaluated as mild, moderate, and severe. Ultrasound of the breast including the axilla was done for all cases and findings were recorded to correlate with clinical features. **Results:** Out of 90 patients studied, the majority was in the age group (21-30) years, (43%) followed by (31-40 years) (40%) respectively. 57(63%) were married and 33(43%) were single. 21 patients had 1 child, 18 had 2 children with 6 of them having no children. Out of 90 patients, only 3 attained menopause, the rest of them have their monthly cycles. 51 patients had cyclical breast pain and 39 had noncyclical breast pain, 39 had pain on the right breast, 36 on left side and bilateral in 15 patients. The pain was mild in 15 patients, moderate in 27 patients and severe in 6 patients. Various USG findings of the affected breast were a normal study in 48 patients followed by duct ectasia and heterogeneous ectogenicity fibroadenosis, small cystic lesion and enlarged axillary

ORIGINAL RESEARCH ARTICLE

Corresponding author*
H. N. Ashikur Rahaman

lymph nodes etc. **Conclusion:** The study results show that majority of patients with breast pain without clinically palpable lump had normal USG study. But it also detects other early changes in the breast tissue. It can be used as a baseline investigation for any breast pain without palpable lump.

©2021, www.medrech.com

I INTRODUCTION

Mastalgia is defined as pain, dull ache, discomfort, or heaviness in the breast. It's commonly experienced by women, presents up to 69% of women, and can affect any age group. The etiology of breast pain is multifactorial. It may be due to normal cyclical changes in hormone levels or due to certain diseases.¹ So, it can be cyclical, non-cyclical with a peak incidence between (35-45) years.² Cyclical hormonal changes are the most common cause of breast pain. Breast cancer is a rare cancer of breast pain according to only 1.2-6.7%.³ Causes of mastalgia without clinical findings of Modularity or lump are still a challenge for diagnosis and management. Ultrasound of the breast is one of the most convenient, painless, and easily available, and less costly investigations that can be done in women complaining of breast pain.⁴ Breast imaging provides reassurance to the women presenting with breast pain without lump according to Wayne state university's study in 2008.³ The cross-sectional study utility of breast imaging in mastalgia by nasreennath. et al⁴. In 2013 showed that Duct ectasia is a major factor in determining the severity of mastalgia with no significance between clinical and nonclinical mastalgia. Hence breast imaging helps to identify the underlying pathology.⁵

II MATERIALS AND METHODS


Patients with complaints of breast pain in the age group 15 to 60 years with clinically no palpable mass attending to the Dept. of Clinical Oncology, Enam Medical College & Hospital, Savar, Dhaka, Bangladesh between January 2020 to December 2020 were included in the study. This was a prospective cross-sectional descriptive study. Detailed history with particular reference to age, duration of symptoms, Size, menstruation, marital status, parity, lactation, nipple discharge, and tenderness are recorded. Pain intensity was evaluated as mild, moderate, and severe. Clinical findings concerning tenderness and quadrants involved are recorded. Then Ultrasound of the affected breast including the axilla was done in all the cases.

III RESULTS

Out of 90 patients studied, the majority was in the age group (21-30) years, (43%) followed by (31-40 years) (40%) respectively. 57(63%) were married and 33(43%) were single. 21 patients had 1 child, 18 had 2 children with 6 of them having no children. Out of 90 patients, only 3 attained menopause, the rest of them have their monthly cycles as shown in the table below:

Table 1: Demographic distribution of study patients.

S. No.	Characteristics (age in years)	Number	Percentage
1	15-20	9	10%
2	21-30	39	43%
3	31-40	36	40%
4	41-60	6	6.6%

**Fig 1:** Marital status.**Table 2:** Children status of women.

S. No.	No. of Children	No. Of Patients
1	No Children	6
2	1-Children	21
3	2-Children	18
4	3-Children	9
5	4 And Above	3

In this study out of 90 patients, 51 (56) patients had cyclical breast pain and 39 (34) had non-cyclical breast pain. The pain was on the right side in 39 (34) patients, 36 (40) on the left, and bilateral in 15 (16) patients. The majority of the pain was in right lower outer

quadrant (27) and left upper quadrants (18). The intensity of the pain was mild in 57 (65), moderate in 27 (30), and severe in 6 (6), patients according to the visual analog scoring (VAS) system.

Table 3: Characteristics of breast pain in the study.

S. No.	Characteristics	Number	Percentage
1	Type of Pain		
	Cyclical	51	56
2	Side affected		
	Right	39	34
	Left	36	40
3	Bilateral	15	16
	Severity of Pain		
	Mild	57	63
	Moderate	30	33
	Severe	3	3

In the USG examination of the affected breast and axilla are found normal study in 48 (53%) patients, duct ectasia in 12 (13%), multiple small cystic lesions in 9 (10%), fibroadenosis in 9 (10%) patients. Heterogenous

ectagenecity with probe tenderness in 4 (4%), Hypoechoic demarcated nodules 4(4%), and fibroadenosis in 3 (3%) and enlarged axillary lymph node 1 (1%) respectively as shown in the table below:

Table 4: Ultrasound findings of the affected breast.

S. No.	Findings	Numbers of Patients	Percentage
01	Normal	48	53
02	Duct Ectasia	12	13
03	Multiple Small Cystic Lesions	9	10
04	Fibroadenosis	9	10
05	Heterogonous Ectagenecity With Probe Tenderness	4	4
06	Hypoechoicdemarcated Nodules	4	4
07	Fibroadenosis	3	3
08	Enlarged Axillary Lymph Node	1	1

Those patients with mass lesions were further subjected to FNAC but no malignancy was detected. FNAC of axillary lymph nodes showed non-specific reactive Lymphadenitis.

IV DISCUSSION

Risk increases by almost 3% for each year older at menopause thus women who have attained menopause at 55 years rather than 45 years, have approximately 30% higher risk. In this study, the majority of the women reached menopause after the age of 45 years. The risk of developing breast cancer increased in both pre-and post-menopausal patients who had early onset of menarche and late menopause possibly due to the increase in the duration of hormonal exposure¹⁵. In our study, more patients were postmenopausal 64% (16 patients) than premenopausal 36% (9 patients). Similar profile can be seen in Karlsson N YA et al series.^{6,7} In Sandhu DS et al⁸ study 44.27% of the patients were premenopausal and 55.76%. Karlsson YA et al reported 41% premenopausal and 59% postmenopausal. Compared to the west the percentage of premenopausal patients is less in this study. Early age at first full-term pregnancy is inversely related to breast cancer risk.⁹ This association perhaps reflects either a pregnancy-induced maturation of mammary cells and thus making them less susceptible to carcinogenic transformation or a long-lasting hormonal change or both.¹⁰ In this study, it was found that the majority of the women

were at an age younger than 30 during their first pregnancy. However, information regarding the age at last pregnancy was not clear. Late age at last full-term pregnancy also is associated with a higher risk of breast cancer, but not in all studies.¹¹ High parity has generally been associated with low breast cancer risk in previous epidemiological studies. Nulliparity was associated with an overall increased risk of breast cancer. In our study, it was found that many women presented with breast cancer despite high parity.¹²

V CONCLUSION

The study results show that the majority of patients with breast pain without clinically palpable lump had normal USG study. But it also detects other early changes in the breast tissue. It can be used as a baseline investigation for any breast pain without a palpable lump.

REFERENCES:

1. Kumar HC, Khadri SI, Kemparaj T, et al. Benign breast diseases: our institutional experience. JEMDS2014; 3(62):13637-44.
2. Memon A, Parveen S, Sangrarasi AK, et al. Changing pattern of benign breast lumps in young females. World J Med Sci 2007; 2(1):21-4.
3. Morris KT, Pommier RF, Morris A, et al. Usefulness of the triple test score for

- palpable breast masses. Arch Surg2001; 136(9):1008-13.
 4. Johnsen C. Breast disease. A clinical study with special reference to diagnostic procedures. Acta Chir Scand Suppl 1975; 454:1-108.
 5. Bassett LW, Ysrael M, Gold RH, et al. Usefulness of mammography and sonography in women less than 35 years of age. Radiology1991; 180(3):831-5.
 6. Ertem G, Kocer A. Breast examination among nurses and midwives in Odemis health district in Turkey. Indian J Cancer2009; 46(3):208-13.
 7. hekwaba FN. Benign breast disease in Nigerian women: a study of 657 patients. J R Coll Surg Edinb1994; 39(5):280-3.
 8. Greenberg R, Skornick Y, Kaplan O. Management of breast fibroadenomas. J Gen Intern Med1998; 13(9):640-5.
 9. Nadia AG. Benign breast disorders-a histopathological studying Tikrit city. Diyala J Med2012; 3(1):27-35.
 10. Gupta A, Gupta AK, Goyal R, et al. A study of clinical profile of benign breast diseases presenting at a tertiary care center in central India. Scholar J Appl Med Sci2015; 3(2C):695-700.
 11. Chaudhary IA, Qureshi SK, Rasul S, et al. Pattern of benign breast diseases. J Surg Pak 2003; 8:5-7.
 12. Sharma A, Chanchlani R. A study of the spectrum of benign breast disease in a tertiary care institute of central India. J Evi Based Med Healthcare2015; 2(5):551-5.
-